

## TEXAS DEPARTMENT OF STATE HEALTH SERVICES

P.O. Box 149347 Austin, Texas 78714-9347 1-888-963-7111 TTY: 1-800-735-2989 www.dshs.state.tx.us

DAVID L. LAKEY, M.D. COMMISSIONER

## \*\*INFLUENZA HEALTH ALERT\*\*

December 20, 2013

## Dear Colleague:

Statewide influenza-like illness (ILI) activity continues to increase and is above baseline levels. ILI intensity is high in Texas, and **influenza is now widespread**. All Texas regions have reported laboratory confirmed influenza. Over 90% of positive influenza tests reported from Texas laboratories have been typed as influenza A. Of those influenza A viruses that have been subtyped, 90% have been the 2009 pandemic H1N1 subtype. This subtype of influenza is included in this season's influenza vaccine.

No novel influenza cases have been reported in Texas.

No antiviral resistant influenza strains have been reported in Texas.

Encourage patients to get vaccinated for influenza.

Clinicians should consider antivirals even if the Rapid Influenza Diagnostic Test is negative.

**Background:** Influenza viruses can be spread by large respiratory droplets generated when an infected person coughs or sneezes in close proximity to an uninfected person. Symptoms can include fever, dry cough, sore throat, headache, body aches, fatigue, and nasal congestion. Among children, otitis media, nausea, vomiting, and diarrhea are common. Most people generally recover from illness in 1-2 weeks, but some people develop complications and may die from influenza. The highest rates of influenza infection occur among children; however, the risks for serious health problems, hospitalizations, and deaths from influenza are higher among people 65 years of age or older, very young children, and people of any age who have medical conditions that place them at increased risk for complications from influenza (see Treatment).

<u>Vaccination:</u> Everyone who is at least 6 months of age should get a flu vaccine this season. It is not too late for vaccination. There are several flu vaccine options available for the 2013-2014 flu season. All these vaccines contain the currently circulating H1N1 strain. DSHS does not recommend one flu vaccine over another, although there are special indications for some (such as a high-dose inactivated trivalent vaccine approved for persons age 65 years and older).

The 2013-2014 trivalent influenza vaccine is made from the following three viruses:

- An A/California/7/2009 (H1N1)pdm09-like virus
- An A(H3N2) virus antigenically like the cell-propagated prototype virus A/Victoria/361/2011
- A B/Massachusetts/2/2012-like virus

The quadrivalent vaccine contains the above three viruses and a B/Brisbane/60/2008-like virus.

Rapid Lab Tests: Rapid Influenza Diagnostic Tests (RIDTs) can be useful to identify influenza virus infection, but false negative test results are common during influenza season. Clinicians should be aware that a negative RIDT result does NOT exclude a diagnosis of influenza in a patient with suspected influenza. When there is clinical suspicion of influenza and antiviral treatment is indicated, antiviral treatment should be started as soon as possible, even if the result of the RIDT is negative, without waiting for results of additional influenza testing.

<u>Treatment</u>: Oseltamivir and zanamivir are chemically related antiviral medications known as neuraminidase inhibitors that have activity against both influenza A and B viruses. Early antiviral treatment can shorten the duration of fever and illness symptoms, may reduce the risk of complications and death, and may shorten the duration of hospitalization. Clinical benefit is greatest when antiviral treatment is administered early, especially within 48 hours of influenza illness onset. Decisions about starting antiviral treatment should not wait for laboratory confirmation of influenza.

Antiviral treatment is recommended as early as possible for any patient with confirmed or suspected influenza who

- Is hospitalized.
- Has severe, complicated, or progressive illness.
- Is at higher risk for influenza complications.

Persons at higher risk for influenza complications recommended for antiviral treatment include:

- Children aged younger than 2 years.
- Adults aged 65 years and older.
- Persons with chronic pulmonary (including asthma), cardiovascular (except hypertension alone), renal, hepatic, or hematological disease (including sickle cell disease); metabolic disorders (including diabetes mellitus); or neurologic and neurodevelopment conditions (including disorders of the brain, spinal cord, peripheral nerve, and muscle such as cerebral palsy, epilepsy [seizure disorders], stroke, intellectual disability [mental retardation], moderate to severe developmental delay, muscular dystrophy, or spinal cord injury).
- Persons with immunosuppression, including that caused by medications or by HIV infection.
- Women who are pregnant or postpartum (within 2 weeks after delivery).
- Persons aged younger than 19 years who are receiving long-term aspirin therapy.
- American Indians/Alaska Natives.
- Persons who are morbidly obese (i.e., body-mass index is equal to or greater than 40).
- Residents of nursing homes and other chronic-care facilities.

Clinical judgment, on the basis of the patient's disease severity and progression, age, underlying medical conditions, likelihood of influenza, and time since onset of symptoms, is important when making antiviral treatment decisions for high-risk outpatients.

Additional details regarding antiviral treatment can be found at <a href="http://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm">http://www.cdc.gov/flu/professionals/antivirals/summary-clinicians.htm</a>.

<u>Disease Reporting Requirements/Statute:</u> Several Texas laws (<u>Health & Safety Code</u>, <u>Chapters 81, 84, and 87</u>) require specific information regarding notifiable conditions to be provided to DSHS. Health care providers, hospitals, laboratories, schools, childcare facilities and

INFLUENZA ALERT December 20, 2013 Page 3

others are required to report patients who are suspected of having a notifiable condition (<u>Chapter 97, Title 25, Texas Administrative Code</u>).

In Texas, influenza-associated pediatric mortality is required to be reported within one work day. Clusters or outbreaks of any disease, including influenza, should be reported immediately. Reports of influenza-associated pediatric mortality and influenza or influenza-like illness outbreak should be made to your local health department or to 1-800-705-8868.

David Lakey, M.D.

Commissioner

Texas Department of State Health Services